Aandoening:
Stage/severity:
Pre/Post operation:

Best imaginable health state

100	Periodontal disease (gingivitis)
99	Acute bronchitis (one episode of 2 weeks in an otherwise healthy year)
98	Post living donor renal transplant: Patients with end-stage renal disease
97	
96	Post thoracotomy: Patients with aortic valve insufficiency, or stenosis
95	Post: Patients with resectable lung carcinoma
94	
93	
92	Pre: Mild/Moderate stable angina pectoris
91	
90	
89	Infertility
88	
87	Post: Women with resectable mammacarcinoma, too large for breast saving
86	
85	
84	Pre: Patients with non-complicated collum femoris fracture
83	
82	Pre: Patients with aortic valve insufficiency, or stenosis
81	
80	
79	Rheumatoid arthritis
78	Pre: Patients with arrythmias (WPW, AVNRT, PV, Flut)
77	Pre: Women with resectable mammacarcinoma
76	
75	
74	
73	
72	
71	Pre: Patients with resectable ovarium carcinoma
70	
69	
68	Pre: Patients with cholangeocarcinoma
67	
66	
65	
64	
63	Deafness
62	
61	
60	
59	
58	
57	Blindness
56	Pre: Patients with resectable lung carcinoma
55	Pre: Patients with high-grade glioma WHO Grade 3/4
54	
53	
52	
51	
50	
49	Down's syndrome
48	
47	
46	
45	
44	
43	Pre: End-stage renal disease
42	Pre: Patients with non-complicated humerus fracture
41	
40	
39	
38	
37	Pre: Stage V chronic kidney disease untreated and severe anemia due to diabetes mellitus or hypertension
36	
35	
34	
33	
32	
31	
30	
29	
28	
27	
26	
25	
24	Unipolar major depression
23	
22	
21	
20	
19	
18	
17	
16	
15	
14	Quadriplegia
13	
12	
11	
10	
9	
8	
7	
6	Dementia
5	
4	
3	High level spina bifida aperta (L2 or higher), permanent stage
2	Active psychosis
1	Post: Colorectal cancer, irradically removed or metastatic carcinoma
0	

Worst imaginable health state